



CellGenix<sup>®</sup> GMP Recombinant Human Interleukin-4 (rh IL-4) Order No.: 1003-050 (50 µg), 1003-1000 (1000 µg)

# **Product Characteristics**

Source	E. coli
Description	Human Interleukin-4, accession # P05112, His25-Ser153 N-terminal Met Molecular mass 15.1 kDa
Formulation	Lyophilized from a 0.2 $\mu$ m-filtered solution containing 1.5 mM potassium phosphate, 8.1 mM sodium phosphate, 2.7 mM potassium chloride and 137 mM sodium chloride, pH 7.5
Intended use	For clinical ex vivo use. Not intended for human in vivo application.

## **Quality Parameters**

ldentity	Confirmed by DNA-sequencing of the expression plasmid in end-of-production cells and N-terminal sequencing of the final product
Activity	8 - 14 x 10 <sup>6</sup> IU/mg <sup>1</sup> calibrated against the NIBSC #88/656
	Measured in a cell proliferation assay using an IL-4-dependent cell line, TF1
Purity	$\geq$ 97 %, as determined by SDS-PAGE (under reducing and non-reducing conditions, visualized by silver staining)
Product-related proteins	$\leq$ 10 % oligomers as determined by SDS-PAGE (under reducing and non-reducing conditions, visualized by silver staining)
Host-cell DNA	$\leq$ 1.0 ng/mg, as determined with the Threshold <sup>®</sup> system <sup>2</sup>
Host-cell protein	$\leq$ 1.0 µg/mg, as determined by ELISA
Endotoxin	$\leq$ 50 EU/mg, as determined by LAL gel clot test according to Ph. Eur. and USP
Sterility	Sterility test according to Ph. Eur. and USP of the vialed product (direct inoculation)
Mass per vial	<b>50 μg:</b> 43 – 57 μg, <b>1000 μg:</b> 900 – 1100 μg as determined by spectrophotometrical measurement
Animal-derived component-free	<b>ADCF Level 2:</b> The final product contains neither animal- nor human-derived materials. ADCF Level 2 cytokines are produced in our dedicated animal-free facility. No animal-derived components are used throughout the complete production process. All ADCF Level 2 cytokines are produced in <i>E. coli</i> .

<sup>&</sup>lt;sup>1</sup> USP units are equivalent to International Units per mg (IU/mg)

<sup>&</sup>lt;sup>2</sup> Threshold<sup>®</sup> System is a trademark of Molecular Devices, Inc.



the

### Shipment & Storage

Transport	Ambient temperature. Please refer to Technote (www.cellgenix.com)
Expiry	Minimum 6 months from date of shipping
Storage & Stability	Store lyophilized cytokine at -20°C to -80°C. Store a 250 μg/ml cytokine solution
	<ul> <li>4 weeks at 2°C to 8°C under sterile conditions after reconstitution. Store in t original container.</li> </ul>

 4 months at -20°C to -80°C under sterile conditions after reconstitution. Store in 60 μl aliquots in polypropylene cryogenic vials.

Avoid repeated freeze/thaw cycles.

### Handling Instructions

ReconstitutionRecommended in sterile water to a final concentration of 250 μg/ml for 50 μg vials or<br/>500 μg/ml for 1000 μg vials.DilutionRecommended in CellGenix® serum-free media. For dilution with protein free medium,<br/>a carrier protein (0.1-1 % albumin or 1-10 % appropriate serum) has to be included.<br/>Failure to dilute product according to these instructions may result in loss of activity.

## **Quality Statement**

This product is manufactured, tested and released in compliance with the relevant GMP-guidelines. USP chapter <1043> has been considered in the design of this product. This product is compliant with Ph. Eur. general chapter 5.2.12.

HQ: CellGenix GmbH | Am Flughafen 16 | D - 79108 Freiburg | GERMANY | Phone +49 761 888 89-0 | Fax +49 761 888 89-830 US: CellGenix Inc. | One New Hampshire Avenue, Suite 125 | Portsmouth, NH 03801 | USA | Phone +1 603 373 0408 | Fax +1 603 373 8104 www.cellgenix.com | info@cellgenix.com

