

Data Sheet

CellGenix® Recombinant Human Interleukin-21 (rh IL-21)

Preclinical Grade - Order No.: 1419-010 (10 µg), 1419-050 (50 µg)

Product Characteristics

Source	<i>E. coli</i>
Description	Human Interleukin-21, accession # Q9HBE4, Gln25-Ser155, N-terminal Met Molecular mass 15.4 kDa
Formulation	Lyophilized from a 0.2 µm-filtered protein solution containing 15 mM sodium acetate and 45 mM sodium chloride, pH 5.5.
Intended use	For preclinical <i>ex vivo</i> use. Not intended for therapeutic use.

Quality Parameters

Activity	≥ 0.5 x 10 ⁶ U/mg calibrated against the internal standard "IL-21 RS1" Measured in a cell proliferation assay using an IL-21-dependent cell line, B9
Purity	≥ 95 %, as determined by SDS-PAGE (under reducing and non-reducing conditions, visualized by Coomassie staining)
Endotoxin	< 25 EU/mg, as determined by LAL gel clot test
Sterility	Sterility test of the vial product (direct inoculation)
Mass per vial	1419-010: 10 µg, 1419-050: 50 µg
Animal-derived component-free	ADCF Level 2: The final product contains neither animal- nor human-derived materials. ADCF Level 2 cytokines are produced in our dedicated animal-free facility. No animal-derived components are used throughout the complete production process. All ADCF Level 2 cytokines are produced in <i>E. coli</i> .

Shipment & Storage

Transport	Ambient temperature. Please refer to Technote (www.cellgenix.com)
Expiry	3 years from date of shipment
Storage & Stability	<p>Store lyophilized cytokine at -20°C to -80 °C.</p> <p>Store a 100 µg/ml reconstituted cytokine solution:</p> <ul style="list-style-type: none">• 4 weeks at 2°C to 8°C under sterile conditions after reconstitution. Store in the original container.• 4 months at -20°C to -80°C under sterile conditions after reconstitution. Store in 80 µl aliquots in polypropylene cryogenic vials. <p>Avoid repeated freeze/thaw cycles.</p>

Handling Instructions

Reconstitution	Recommended in sterile water to a final concentration of 100 µg/ml (for 10 µg vials) or 250 µg/ml (for 50 µg vials).
Dilution	Recommended in CellGenix® serum-free media. For dilution with protein free medium, a carrier protein (0.1-1 % albumin or 1-10 % appropriate serum) has to be included. Failure to dilute product according to these instructions may result in loss of activity.

Quality Statement

Final manufacturing steps and QC are performed in a GMP facility.